## Student Asthma/Allergy Action Plan

(This Page To Be Completed By Physician)

Student Name:	Date Of Birth:/			
■ Exercise Pre-Treatment: Administer inhaler (2 inhalations) 15-30 minutes prior to exercise. (e.g., PE, recess, etc).				
<ul> <li>□ Albuterol HFA inhaler (Proventil, Ventolin, ProAir)</li> <li>□ Levalbuterol (Xopenex HFA)</li> <li>□ Pirbuterol inhaler (Maxair)</li> </ul>	<ul> <li>Use inhaler with valved holding chamber</li> <li>May carry &amp; self-administer inhaler (MDI)</li> <li>Other:</li> </ul>			
Asthma Treatment	Anaphylaxis Treatment			
Give quick relief medication when student has asthma symptoms, such as coughing, wheezing or tight chest.  Albuterol HFA (Proventil, Ventolin, ProAir) 2 inhalations  Levalbuterol (Xopenex HFA) 2 inhalations  Pirbuterol (Maxair) 2 inhalations  Use inhaler with valved holding chamber  May carry & self-administer inhaler (MDI)  Albuterol inhaled by nebulizer (Proventil, Ventolin, AccuNeb)  Gas mg/3 mL  Levalbuterol inhaled by nebulizer (Xopenex)  Gosal mg/3 mL  Other:  Closely Watch the Student after Giving Quick Relief Medication  If, after 10 minutes:  Symptoms are better, student may return to classroom after notifying parent/guardian  Symptoms are not better, give the treatment again and notify parent/guardian right away  If student continues to get worse, CALL 911 and use the Nebraska Schools' Emergency Response to Life-Threatening Asthma or Systemic Allergic Reactions (Anaphylaxis) Protocol  This student has a medical history of asthma and/or anaphylaxis ar If medications are self-administered, the school staff must be no Additional information: (i.e. asthma triggers, allergens)	Give epinephrine when student has allergy symptoms, such as hives, hard to breathe (chest or neck "sucking in"), lips or fingernails turning blue, or trouble talking (shortness of breath).  □ EpiPen® 0.3 mg □ EpiPen® Jr. 0.15 mg □ Auvi-Q™ 0.3 mg □ Auvi-Q™ 0.15 mg □ Adrenaclick® 0.3 mg □ Adrenaclick® 0.15 mg □ May carry & self-administer epinephrine auto-injector □ Use epinephrine auto-injector immediately upon exposure to known allergen  CALL 911 After Giving Epinephrine & Closely Watch the Student  • Notify parent/guardian immediately • Even if student gets better, the student should be watched for more symptoms of anaphylaxis in an emergency room  • If student does not get better or continues to get worse, use the Nebraska Schools' Emergency Response to Life-Threatening Asthma or Systemic Allergic Reactions (Anaphylaxis) Protocol			
Physician name: (please print)	Phone:			
Physician signature: Date:				
Parent signature:				

\_ Date:\_ Version: 10/13 Page I of 2

Reviewed by school nurse/nurse designee:\_

## Student Asthma/Allergy Action Plan

(This Page To Be Completed By Parent/Guardian)

Student Nan	ne:		Age:	Grade:	
School:		Homeroon	Homeroom Teacher:		
Parent/Guar	dian:		Phone(H)	( <b>W</b> )	
Parent//Guardian:		Phone(H)	(W)		
Alternate Em	ergency Contact:		Phone(H)	(W)	
Known Asth	ma Triggers: Please check the boxes	to identi	fy what can cause an asthm	a episode for your student.	
	Respiratory/viral infections Animals/dander ure/weather—humidity, cold air, etc. lease list:		Odors/fumes/smoke Dust/dust mites Pesticides	<ul><li>☐ Mold/mildew</li><li>☐ Grasses/trees</li><li>☐ Food—please list below</li></ul>	
Known Aller	rgy/Intolerance: Please check those we allergen	hich app	ly and describe what happ	ens when your child eats or comes into	
Peanuts Tree Nuts Fish/shellfish Eggs Soy Wheat Milk Medication Latex Insect stings Other Notice: If your	child has been prescribed epinephrine (sue eeds a special diet to limit or avoid foods,	ich as an	EpiPen®) for an allergy, yo	ou must provide epinephrine at school. If	
Daily Medicines:Please list daily medicines used at home and/or to be given at school.Medicine NameAmount/DoseWhen does it need to given					
l unde	erstand that all medicines to be gi	iven at	school must be provi	ded by the parent/guardian.	
Parent signa	ture:			Date:	
Reviewed by school nurse/nurse designee:				Date:	

Page 2 of 2

Version: 10/13